

# Phylogeography of lions (*Panthera leo* ssp.) reveals three distinct taxa and a late Pleistocene reduction in genetic diversity

ROSS BARNETT,\* BETH SHAPIRO,† IAN BARNES,‡ SIMON Y. W. HO,§ JOACHIM BURGER,¶ NOBUYUKI YAMAGUCHI,\*\* THOMAS F. G. HIGHAM,†† H. TODD WHEELER,‡‡ WILFRIED ROSENDAHL,§§ ANDREI V. SHER,¶¶ †††† MARINA SOTNIKOVA,\*\*\* TATIANA KUZNETSOVA,††† GENNADY F. BARYSHNIKOV,††† LARRY D. MARTIN,§§§ C. RICHARD HARINGTON,¶¶¶ JAMES A. BURNS\*\*\*\* and ALAN COOPER††††

\*Department of Zoology, University of Oxford, Oxford OX1 3PS, UK, †Department of Biology, Pennsylvania State University, University Park, PA 16801, USA, ‡Department of Biology, University College London, London WC1E 6BT, UK, §Centre for Macroevolution and Macroecology, School of Botany and Zoology, Australian National University, Canberra, ACT 0200, Australia, ¶Institute of Anthropology, Mainz University, 55099 Mainz, Germany, \*\*Department of Biological and Environmental Sciences, University of Qatar, PO Box 2713, Doha, Qatar, ††Research Laboratory for Archaeology, University of Oxford, Oxford OX1 3QJ, UK, ‡‡George C. Page Museum of La Brea Discoveries, Los Angeles, CA 90036, USA, §§Reiss-Engelhorn-Museen, D-68159 Mannheim, Germany, ¶¶Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, 119071 Moscow, Russia, \*\*\*Geological Institute, Russian Academy of Sciences, 119017 Moscow, Russia, †††Palaeontological Department, Geological Faculty, Moscow State University, 119899 Moscow, Russia, ††††Zoological Institute, Russian Academy of Sciences, 199034 St Petersburg, Russia, §§§Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, KS 66045, USA, ¶¶¶Canadian Museum of Nature (Palaeobiology), Ottawa, ON, Canada K1P 6P4, \*\*\*\*Quaternary Palaeontology, Royal Alberta Museum, Edmonton, AB, Canada T5N 0M6, ††††School of Earth and Environmental Sciences, University of Adelaide, Adelaide, SA 5005, Australia

## Abstract

Lions were the most widespread carnivores in the late Pleistocene, ranging from southern Africa to the southern USA, but little is known about the evolutionary relationships among these Pleistocene populations or the dynamics that led to their extinction. Using ancient DNA techniques, we obtained mitochondrial sequences from 52 individuals sampled across the present and former range of lions. Phylogenetic analysis revealed three distinct clusters: (i) modern lions, *Panthera leo*; (ii) extinct Pleistocene cave lions, which formed a homogeneous population extending from Europe across Beringia (Siberia, Alaska and western Canada); and (iii) extinct American lions, which formed a separate population south of the Pleistocene ice sheets. The American lion appears to have become genetically isolated around 340 000 years ago, despite the apparent lack of significant barriers to gene flow with Beringian populations through much of the late Pleistocene. We found potential evidence of a severe population bottleneck in the cave lion during the previous interstadial, sometime after 48 000 years, adding to evidence from bison, mammoths, horses and brown bears that megafaunal populations underwent major genetic alterations throughout the last interstadial, potentially presaging the processes involved in the subsequent end-Pleistocene mass extinctions.

*Keywords:* American lion, ancient DNA, Beringia, cave lion, extinction, megafauna

*Received 3 October 2008; revision received 6 January 2009; accepted 15 January 2009*

## Introduction

Lions currently survive only in parts of Africa and India, yet as recently as the late Pleistocene (LP), large and morphologically diverse forms of lion were distributed across the Northern Hemisphere. Lions survived in Europe,

Correspondence: Ross Barnett, E-mail: drrossbarnett@gmail.com; Alan Cooper, Australian Centre for Ancient DNA, School of Earth and Environmental Sciences, University of Adelaide, Adelaide, SA 5005, Australia. E-mail: alan.cooper@adelaide.edu.au  
††††Deceased.




Fig. 1 Pleistocene lion distribution and sample origin. Colours correspond to commonly cited population subdivisions (e.g. Baryshnikov & Boeskorov 2001). Numbers refer to particular samples used in this analysis (Table 1).

northern Asia, and North America until around 11 000 BP (Fig. 1), when they disappeared as part of the end-Pleistocene megafaunal mass extinctions (Martin & Steadman 1999). Human hunting and/or environmental alteration, and climate changes associated with the last glacial maximum (LGM; 22 000–18 000 BP) and the end of the Pleistocene have variously been implicated (Barnosky *et al.* 2004).

In eastern Beringia, a Pleistocene landmass consisting of present-day Alaska and Canada's Yukon Territory, genetic and palaeontological data from large mammal populations suggest that a key period of change occurred around 40 000–21 000 BP, in the period before the LGM (Barnes *et al.* 2002; Guthrie 2003; Shapiro *et al.* 2004). Due to their worldwide distribution, genetic analyses of Pleistocene lions potentially hold key insights into how megafaunal carnivore populations responded to environmental changes in different regions. Additionally, a better understanding of how and when lion populations went extinct is vital for reconstructing community-level dynamics during the LP, which in turn is fundamental for testing hypotheses about the cause of the end-Pleistocene extinctions.

All Pleistocene and modern lion specimens are assigned to the genus *Panthera*, but there is little consensus as to the

number of distinct species or the extent of overlap of their distributions (for a comprehensive overview of the taxonomy of pantherine cats, please refer to Burger *et al.* 2004). Morphological identification is complicated by the presence of shared primitive features (e.g. Sotnikova & Nikolskiy 2006), although characteristics such as body size or the presence/absence of a mane in males have been used to suggest taxonomic distinctiveness between geographical regions (e.g. Kurtén 1968; Yamaguchi *et al.* 2004; Sotnikova, Nikolskiy 2006). Pleistocene lions in Europe are usually referred to as *spelaea*, while North American lions are given the designation *atrox*. However, the status of these groups as species or subspecies, and their relationships to each other and to extant *Panthera* remains unclear (e.g. Simpson 1941; Kurtén 1985; Harington 1971; Vereshchagin 1971; Kurtén & Anderson 1980; Herrington 1986; Groiss 1996; Turner & Antón 1997; Baryshnikov & Boeskorov 2001; Burger *et al.* 2004; Sotnikova, Nikolskiy 2006).

In this study, the first to examine intraspecific diversity in an extinct Pleistocene felid, we investigate the relationships among the putative lion subspecies. This was carried out by obtaining mitochondrial DNA sequences from modern and ancient lions. The evolutionary and

demographic history of the sequences is inferred using phylogenetic analysis, and the results are interpreted in the context of environmental changes in the late Pleistocene.

### Materials and methods

More than 200 cortical bone samples were obtained from museum specimens of modern, Pleistocene, and American lions (Table S1, Supporting information). Specimens ranged in age from modern to > 62 000 years. To prevent duplicate sampling within a site, either the same skeletal element or samples associated with widely separated radiocarbon dates were chosen. DNA extraction was performed as described in Barnett *et al.* (2006) in dedicated ancient DNA (aDNA) facilities at the University of Oxford. Four European lions were also extracted and amplified at Mainz University under similarly stringent conditions (Burger *et al.* 2004). One of the cave lions from Mainz (sample 8 in Table 1) was previously analysed for cytochrome *b* in an earlier study showing the phylogenetic position of the cave lion among the *Panthera* cats (Burger *et al.* 2004).

A ~215-bp fragment of the mitochondrial hypervariable region 1 (HVR1) and a 143-bp fragment of *ATP8* were polymerase chain reaction (PCR) amplified, and cloned, purified, and sequenced following Barnett *et al.* (2006). Strict aDNA protocols were followed, including the use of multiple PCRs and negative controls, overlapping fragments and independent replication. Nuclear mitochondrial inserts, which are known to be widespread among felids (e.g. Kim *et al.* 2006), were identified and excluded from the analysis. All specimens possessing unique haplotypes, along with a subset of those with shared haplotypes, were amplified several times and cloned to check for the presence of contamination (details are given in Supporting Information: supporting text, Tables S1–S3).

Thirty-three samples were submitted to the Oxford Radiocarbon Accelerator Unit (University of Oxford) for radiocarbon analysis. Analyses were performed using 0.2 g of bone taken from a site adjacent to the sample used for DNA extraction. An additional sample (sample 8 in Table 1) was radiocarbon dated at the Leibniz laboratory of the University of Kiel. For all samples, total bone collagen was extracted, graphitized, and dated by accelerator mass spectrometry. Dates are presented as uncalibrated radiocarbon values. Two of the oldest specimens (samples 31 and 34 in Table 1) were dated twice, to test the reproducibility of dates in the > 50 000-year time frame.

Phylogenetic analyses were performed on the HVR1 and *ATP8* data sets separately, due to the differing number of specimens that yielded sequences from the two regions (Supporting Information: supporting text, Table S3). Median-joining networks were produced for both data sets using the program Network version 4.1.0.3 (Bandelt *et al.* 1999).

Bayesian Markov chain Monte Carlo (MCMC) analyses were then performed using BEAST (Drummond & Rambaut 2007), first for HVR1 and *ATP8* data sets separately, and then for a smaller, combined analysis consisting of only those samples from which both sequence fragments could be amplified and which were associated with finite radiocarbon dates. For all three data sets, comparison of Akaike information criterion scores suggested the HKY85 model of nucleotide substitution. Each BEAST analysis assumed this substitution model as well as a constant population size and a strict molecular clock calibrated using the age of the split between the *spelaea* and *leo* groups (with a normal prior mean of 550 000 years, standard deviation 25 000 years), based on the first appearance of *Panthera leo fossilis* (Garcia Garcia 2001; Burger *et al.* 2004). Demographic and evolutionary model parameters were then estimated simultaneously along with the phylogeny, with samples drawn from the posterior every 5000 MCMC steps over a total of 5 000 000 steps. The first 500 000 steps were discarded as burn-in. Acceptable mixing and convergence to stationarity were checked using the program Tracer version 1.4 (Rambaut & Drummond 2007).

### Results

DNA was successfully amplified from 38 LP and 14 modern lions (Fig. 1, Table S3, Supporting Information). The repeated extractions, amplifications, and cloning experiments yielded consistent results and negative controls at the extraction and PCR stages were uniformly clean. Collagen levels in the samples submitted for radiocarbon analysis were consistently high, indicating good biochemical preservation. The specimens older than 50 000 years that were dated twice gave results within one standard deviation (Supporting information). The sequences contained 48 parsimony-informative sites, distinguishing 24 distinct haplotypes. These are subdivided into three well-supported clusters in both Bayesian (Fig. 2) and median-joining network analyses (Fig. 3): modern lion (*Panthera leo*, from Africa and southwest Asia), cave lion (*spelaea*, from Europe, Siberia, Alaska, and the Yukon) and American lion (*atrox*, from south of the LP glacial ice sheets in North America). All late Pleistocene lion samples produced sequences that grouped strongly with modern lion data (Jae-Heup *et al.* 2001; Barnett *et al.* 2006), rejecting any postulated link between *atrox* and jaguar (*Panthera onca*) (Simpson 1941; Christiansen 2008) or *spelaea* and tiger (*Panthera tigris*) (Herrington 1986; Groiss 1996).

Previous work has shown nuclear translocations of mitochondrial DNA (numts) to be common in cats (Lopez *et al.* 1994; Lopez *et al.* 1996; Cracraft *et al.* 1998; Kim *et al.* 2006). A possible numt of the lion HVR1 was found in only one of the 15 modern lions (sample 50 in Table 1). This numt was similar to that previously reported for other


**Table 1** List of samples used in this study along with museum accession numbers, sample provenance, haplotype assignment, radiocarbon dates and whether the extract was also used in an earlier study

| Sample number | Control region haplotype | Extract | Element     | Museum accession no. | Origin  | Uncalibrated radiocarbon date | OxAU accession no.             | Partial sequence also in Barnett <i>et al.</i> (2006) |
|---------------|--------------------------|---------|-------------|----------------------|---|-------------------------------|--------------------------------|---|
| 1             | A                        | IB133   | L. Humerus  | CMN 47294            | Gold Run Ck, Yukon, Canada                      | 12 640 ± 75                   | OxA-10083                      | YES   |
| 2             | A                        | RB112   | Canine Root | CMN 43789            | Caribou Ck, Yukon, Canada, 1985                 | n/a                           | n/a                            | n/a   |
| 3             | A                        | RB74    | Phalanx     | FAM69073             | Fairbanks Ck, Alaska, USA, 1949                 | n/a                           | n/a                            | n/a   |
| 4             | A                        | RB75    | Tibia       | FAM69167             | Ester Ck, Alaska, USA, 1938                     | 12 090 ± 80                   | OxA-13451                      | YES   |
| 5             | B                        | IB134   | Ramus       | FAM 69016            | Gold Hill, Alaska, USA, 1952                    | 18 240 ± 90                   | OxA-10084                      | n/a   |
| 6             | B                        | IB136   | R. Femur    | CMN 35609            | Hunker Ck, Dawson, Canada, 1978                 | 15 550 ± 90                   | OxA-10086                      | n/a   |
| 7             | B                        | IB138   | Humerus     | FAM A-780-1435       | Fairbanks Ck, Alaska, USA, 1955                 | 11 925 ± 70                   | OxA-10080                      | n/a   |
| 8             | B                        | MZ124   | Pelvis      | MKU                  | Tischhofer Höhle, Austria                       | 31 890 ± 300                  | KIA 16510                      | n/a   |
| 9             | B                        | RB18    | Humerus     | FAM 69105            | Engineer Ck, Alaska, USA, 1941                  | n/a                           | n/a                            | n/a   |
| 10            | B                        | RB36    | Radius      | PIN 3020-350         | Alazeya River, Russia, 1972                     | 13 770 ± 55                   | OxA-13835                      | n/a   |
| 11            | B                        | RB39    | Ulna        | IEM 153-003          | Arga-Yurekh River, Russia, 1982                 | 12 525 ± 50                   | OxA-13833                      | n/a   |
| 12            | B                        | RB41    | Calcaneus   | GIN 772-95/1341      | Duvanny Yar, Loc.1341, Russia, 1964             | 46 200 ± 1500                 | OxA-13024                      | YES   |
| 13            | B                        | RB45    | Radius      | GIN xDYa-84          | Duvanny Yar, Russia, 1984                       | 28 720 ± 160                  | OxA-12981                      | YES   |
| 14            | B                        | RB47    | Ulna        | PIN 3020-73          | Beryozovka River, Russia, 1970                  | 27 950 ± 140                  | OxA-13831                      | n/a   |
| 15            | B                        | RB72    | Humerus     | FAM69101             | Fairbanks Ck, Alaska, USA, 1950                 | n/a                           | n/a                            | n/a   |
| 16            | B                        | RB73    | Humerus     | FAM69104             | Lower Gold Stream, Alaska, USA, 1939            | n/a                           | n/a                            | n/a   |
| 17            | B                        | RB77    | Ulna        | FAM69138             | Fairbanks Ck, Alaska, USA, 1951                 | 17 890 ± 100                  | OxA-13452                      | n/a   |
| 18            | B                        | RB91    | Femur       | ZIN 29421(2)         | Lena River, Russia                              | 12 450 ± 60                   | OxA-12901                      | n/a   |
| 19            | B                        | RB96    | Bone        | UAF IK-01-409        | North Slope, Alaska, USA                        | 12 630 ± 60                   | OxA-13473                      | n/a   |
| 20            | B                        | RB157   | Bone        | UB J.S.951           | Ural Mountains, Russia                          | ~13 000                       | Date supplied by A. Hufthammer | n/a   |
| 21            | C                        | RB17    | Ulna        | FAM 69139            | Fairbanks Ck, Alaska, USA, 1954                 | 16 005 ± 65                   | OxA-13834                      | n/a   |
| 22            | C                        | RB24    | Ramus       | FAM 30757            | Banner Ck, Alaska, USA, 1938                    | 15 975 ± 65                   | OxA-13832                      | n/a   |
| 23            | D                        | RB60    | Tibia       | VIE GS-27            | Gamssulzen, Höhle, Austria                      | 49 900 ± 1500                 | OxA-13110                      | YES   |
| 24            | E                        | MZ477b  | Femur       | UEN                  | Gremsdorf, Germany                              | 28 310 ± 50                   | OxA-14862                      | n/a   |
| 25            | F                        | MZ475   | Phalanx     | UMB                  | Zoolithenhöhle, Germany                         | 47 600 ± 900                  | OxA-14863                      | n/a   |
| 26            | G                        | RB43    | Mandible    | IPF AR(DYA)2002      | Duvanny Yar, Russia, 2002                       | > 53 200                      | OxA-13022                      | n/a   |
| 27            | G                        | RB64    | Calcaneus   | STU 9995.2           | Sibyllen Höhle, Germany                         | > 48 100                      | OxA-15354                      | YES   |
| 28            | H                        | RB115   | Tibia       | IEM bl-0418-1        | Bolshoi Liakhovsky Island, Russia, shore 1998   | > 62 100                      | OxA-13837                      | n/a   |
| 29            | H                        | RB116   | Bone        | IEM bl-0192-1        | Bolshoi Liakhovsky Island, Russia, in situ 1998 | > 60 700                      | OxA-13836                      | n/a   |
| 30            | H                        | RB42    | Femur       | PIN 3915-121         | Khromskaya Guba, Khaptashinskiy, Russia, 1978   | > 50 600                      | OxA-13023                      | n/a   |
| 31            | I                        | RB38    | Calcaneus   | PIN 2998-217         | Chukochya, Loc. N 27, Kolyma, Russia, 1969      | 55 700 ± 3000                 | OxA-13025                      | n/a   |
|               |                          |         |             |                      |   | 54 600 ± 1700                 | OxA-13475                      |   |
| 32            | J                        | RB44    | Radius      | PIN 3916-162         | Krestovka River, Loc.6, Russia, 1979            | 54 100 ± 1800                 | OxA-13830                      | n/a   |
| 33            | K                        | RB46    | Femur       | PIN 3752-24A         | Duvanny Yar, Russia, 1976                       | > 61 500                      | OxA-13829                      | n/a   |
| 34            | L                        | RB48    | Calcaneus   | GIN KON.345          | Alazeya River, Russia, 1950s                    | 55 300 ± 2500                 | OxA-13021                      | n/a   |
|               |                          |         |             |                      |   | 58 200 ± 3500                 | OxA-13474                      |   |

Table 1 Continued

| Sample number | Control region haplotype | Extract     | Element            | Museum accession no.   | Origin                                | Uncalibrated radiocarbon date | ORAU accession no. | Partial sequence also in Barnett <i>et al.</i> (2006) |
|---------------|--------------------------|-------------|--------------------|------------------------|---------------------------------------|-------------------------------|--------------------|---|
| 35            | M                        | IB174/RB153 | Humerus            | KU44409                | Natural Trap Cave, Wyoming, USA       | 24 080 ± 170                  | OxA-10078          | n/a   |
| 36            | N                        | IB177/RB154 | Humerus            | KU44000                | Natural Trap Cave, Wyoming, USA       | n/a                           | n/a                | n/a   |
| 37            | O                        | RB113       | R. Mandible        | EDM P98.5.404          | Consolidated pit 48, Edmonton, Canada | 28 940 ± 240                  | OxA-13453          | n/a   |
| 38            | P                        | RB95        | L. Metatarsal II   | EDM P89.13.546         | Consolidated pit 48, Edmonton, Canada | 11 355 ± 55                   | OxA-12900          | n/a   |
| 39            | Q                        | RB206       | Mandible           | STK A58:5287           | Barbary, 1831                         | n/a                           | n/a                | YES   |
| 40            | R                        | IB 4,RB243  | Drilled Bone       | PAR 1962-2854          | Iran                                  | n/a                           | n/a                | YES   |
| 41            | S                        | IB 27       | Drilled Bone       | PAR cA-1892            | Senegal                               | n/a                           | n/a                | n/a   |
| 42            | T                        | IB43        | Tissue             | EDI                    | India                                 | n/a                           | n/a                | YES   |
| 43            | T                        | PLP         | Bone               | MAN                    | India                                 | n/a                           | n/a                | n/a   |
| 44            | U                        | IB 6        | Drilled Bone       | PAR 1995-164           | Sudan                                 | n/a                           | n/a                | YES   |
| 45            | V                        | RB209       | Drilled Bone       | STK A58:1971           | Namibia                               | n/a                           | n/a                | YES   |
| 46            | V                        | PLE-181     | Published Sequence | Jae-Heup <i>et al.</i> | Serengeti, Tanzania                   | n/a                           | n/a                | n/a   |
| 47            | V                        | RB215       | Skull              | BAW 5728               | Zambia                                | n/a                           | n/a                | YES   |
| 48            | V                        | RB216       | Skull              | STK A59:5062           | DRC (NE), 1921                        | n/a                           | n/a                | YES   |
| 49            | V                        | RB217       | Mandible           | STK A59:5066           | DRC (NE), 1921                        | n/a                           | n/a                | YES   |
| 50            | W                        | RB210       | Drilled Bone       | AMA 15904              | South Africa                          | n/a                           | n/a                | YES   |
| 51            | W                        | RB213       | Skull              | BAW 29119              | Botswana                              | n/a                           | n/a                | YES   |
| 52            | W                        | RB214       | Skull              | BAW 63591              | Zimbabwe, 1967                        | n/a                           | n/a                | YES   |
| 53            | X                        | RB208       | Drilled Bone       | AMA 107.1              | Tanzania                              | n/a                           | n/a                | YES   |
| 54            | ?                        | IB135       | Ulna               | FAM 69158              | Cripple Ck Sump, Alaska, USA, 1950    | 53 900 ± 2300                 | OxA-10085          | n/a   |
| 55            | ?                        | IB153       | Bone               | KRA 6857               | Wierchowska, Poland                   | 38 650 ± 600                  | OxA-10087          | n/a   |
| 56            | ?                        | IB81        | Humerus            | FAM69094               | Lower Gold Stream, Alaska USA, 1939   | 12 540 ± 75                   | OxA-10081          | n/a   |
| Leopard       | <i>P. pardus</i>         | IB123       | Tissue             | EDI                    | Saudi Arabia                          | n/a                           | n/a                | n/a   |
| Tiger         | <i>P. tigris</i>         | RB100       | Tissue             | EDI                    | Siberia, Russia                       | n/a                           | n/a                | n/a   |
| Jaguar        | <i>P. onca</i>           | RB52        | Tissue             | EDI                    | South America                         | n/a                           | n/a                | n/a   |

Museum code: American Museum of Natural History (AMNH), Canadian Museum of Nature (CMN), University of Vienna (VIE), Mannheim Museum (MAN), University of Bergen (UB), State Museum of Natural History Stuttgart (STU), University of Alaska (UAF), Palaeontological Institute Moscow (PIN), Russian Academy of Sciences (IEM), Moscow State University (GIN), Zoological Institute St. Petersburg (ZIN), Polish Academy of Sciences (KRA), Royal Alberta Museum (EDM), University of Kansas (KU), Muséum national d'Histoire naturelle (PAR), Amathole Museum (AMA), Natural History Museum of Zimbabwe (BUW), Swedish Museum of Natural History (STK), University of Erlangen-Nuremberg (UEN), Urweltmuseum Bayreuth (UMB), Museum Kufstein (MKU), and the Royal Museum of Scotland (EDI). All sequences have been deposited in GenBank (DQ899900–DQ899945, DQ318552–DQ318562).



**Fig. 2** Evolutionary relationships among lions estimated using Bayesian phylogenetic analysis of the mitochondrial HVR1. Colours and numbers correspond to those in Fig. 1. Nodes with posterior probabilities greater than 0.5 are labelled. Haplotypes and mean uncalibrated radiocarbon dates are given (where available). TMRCA estimated from a combined HVR1 and *ATP8* data set are indicated:  
\*337 000 BP (95% HPD: 194 000–489 000 BP).  
\*\*200 000 BP (95% HPD: 118 000–346 000 BP).

pantherines (Cracraft *et al.* 1998; Kim *et al.* 2006) and was highly divergent from the target mitochondrial locus. A nuclear copy of the *ATP8* gene was also detected in several lion specimens, as well as in leopard and tiger. The *ATP8* numt sequences clustered in a basal position in the phylogeny, and were also distinguished by an adenine insertion in the tiger sequence and a stop codon in the leopard sequence.


Under the assumption of a molecular clock, the substitution rate for the data set of combined HVR1 and *ATP8* sequences was estimated at  $5.9 \times 10^{-8}$  substitutions/site/year, with a 95% highest posterior density (HPD) of  $3.7$ – $8.3 \times 10^{-8}$  substitutions/site/year. Due to the relatively low information content of the sequences, it was not possible to employ a relaxed molecular clock to allow for changes in evolutionary rate over the time period examined (Ho *et al.* 2005). The combined HVR1/*ATP8* analysis resulted in an estimate for the divergence between *spelaea* and *atrox* at

337 000 BP (95% HPD: 194 000–489 000 BP). Estimates for the time to most recent common ancestor (TMRCA) for LP lions are given in Fig. 2. The large temporal range of these estimates reflects both the considerable uncertainty in the primary fossil calibration and the relatively low information content of the sequences.

## Discussion

Three distinct types of lion survived during the LP: the American lion (*atrox*), cave lion (*spelaea*) and the ancestors of the modern lion (*Panthera leo*) (Fig. 1). The phylogenetic topology (Fig. 2) indicates that the mitochondrial lineages of *atrox* and *spelaea* form sister clades.

Network analyses (Fig. 3) produce a similar topology, but place *atrox* inside the basal diversity of *spelaea*, suggesting *atrox* was derived from a Beringian population that dispersed into North America and was subsequently isolated.



**Fig. 3** Graph of uncalibrated radiocarbon date against haplotype sorted by geographical area. Median-joining network analysis of HVR1 and ATP8. Different letters represent different haplotypes, corresponding to those in Table 1. Colours represent area of origin as in Fig. 1. Area of circles are proportional to haplotype frequencies, and the lengths of connecting lines are proportional to the distance between haplotypes (defined as the number of substitutions estimated using Network version 4.1.0.3). Points at which lines intersect without associated circles indicate putative ancestral states.

Molecular date estimates place this initial divergence at 337 000 BP (95% HPD: 194 000–489 000 BP) with a most recent common ancestor of the *atrox* lineage c. 200 000 BP (95% HPD: 118 000–346 000 BP; Fig. 2). This is consistent with palaeontological evidence of lions in eastern Beringia during the Illinoian glacial (c. 300 000–130 000 BP, Harington 1969), and in central North America from the Sangamon interglacial (c. 130 000–120 000 BP). Lions persisted in North America, both north and south of the ice, until the end of the Pleistocene (Harington 1969; Kurtén & Anderson 1980).

The most recent directly radiocarbon dated *atrox* is that from Edmonton, at 11 355 ± 55 BP. (Table 1) This is younger than the lion metatarsal from Jaguar Cave, Idaho, which previously yielded a date of 11 900 ± 130 BP (Kurtén &

Anderson 1980; Stuart 1991). The date of 10 370 ± 150 BP, which has been used previously as the terminal date for *Panthera leo atrox* (Kurtén & Anderson 1980; Stuart 1991) is based indirectly on hearth charcoal from Jaguar Cave. The radiocarbon-dated cave lions from western Beringia reveal a reasonably continuous chronological distribution with a terminal date from the Lena River of 12 450 ± 60 BP (Table 1). The terminal date for eastern Beringia (11 925 ± 70 BP, Fairbanks Creek) is 1500 years older than the widely reported Lost Chicken Creek date of 10 370 ± 160 BP (Kurtén & Anderson 1980; Stuart 1991; Harington 1980), which comes not directly from a lion, but a bison found at the same site. In Eurasia, the youngest dated lion is that reported from Zigeuenerfels, Sigmaringen, Germany at 12 375 ± 50 BP (Stuart & Lister 2007). The widely reported

date from Lathum in the Netherlands ( $10\,670 \pm 160$  BP) (Stuart 1991) has recently been shown to be much older, and therefore invalid (Stuart & Lister 2007).

Modern and LP European cave lions have been shown previously (Burger *et al.* 2004) to be distinct, but this is the first genetic evidence for a grouping of European *spelaea* with lions from Beringia as a single population. Similarly, the genetic separation between LP lions in Beringia and those south of the North American ice sheets is reported for the first time. The reasons for the lack of gene flow within North America remain unclear, as there is no evidence of a barrier between Beringia and central North America during interstadial periods for other taxa such as horse and bison (Burns 1996; Shapiro *et al.* 2004; Weinstock *et al.* 2005). *Atrox* is also present in Edmonton, Alberta, within the region postulated to be an ice-free corridor at various times in the late Pleistocene (Burns 1996; Wilson 1996) both before and after the LGM (Fig. 1, samples 37 and 38 in Table 1). While this indicates that the region was habitable by lions during interstadial periods, it is possible that currently unidentified ecological barriers prevented movement between the Yukon and southern Canada, such as the intermittent presence of forested regions (Wilson 1996) suggested to have been a barrier to the dispersal of bison following the LGM (Shapiro *et al.* 2004). Alternatively, some form of interspecific competition may have existed between *atrox* and *spelaea*, preventing interchange between the areas, as it seems unlikely that environmental or ecological barriers operated over the entire 200 000-year period of separation. Interestingly, there is also no evidence for genetic interchange between cave lions and modern lions, despite a long-term contact zone in the Near East (Fig. 1; Kurtén 1968; Burger *et al.* 2004). These observations are consistent with the suggestion that the three forms may represent separate species, rather than subspecies (Sotnikova, Nikolskiy 2006). However, pronounced mitochondrial phylogeographical structure has also been observed in Pleistocene eastern Beringian brown bear populations, albeit for much shorter time periods (Barnes *et al.* 2002). To fully resolve the species status of the three lion forms, it will be necessary to include both nuclear and morphological data.

The clear genetic distinction between the *atrox* and *spelaea* groups in North America is in sharp contrast to the paucity of genetic divisions within *spelaea* across Europe, Asia, and Alaska/Yukon, suggesting genetic interchange across an immense geographical range. This is particularly apparent after *c.* 46 000–48 000 BP when, despite the small sample size, we observe a dramatic and widespread decline in genetic diversity within *spelaea* across western Beringia and Europe (Figs 2 and 3). Before 46 000 BP, nine diverse mitochondrial haplotypes (B, D, F–L; Figs 2 and 3) are observed from 12 specimens. In contrast, all 18 *spelaea* specimens sampled across the entire range of *spelaea* after this point form part of a closely related star-shaped radiation, consistent with a

rapid population expansion, of four derived haplotypes (A, B, C, and E; Figs 2 and 3) centred around B, the only haplotype shared with the earlier time period. This marked decline in diversity may simply reflect genetic drift, combined with the relatively sparse temporal–geographical sampling. However, this seems somewhat unlikely given the pattern of loss, where all seven known haplotypes from outside the star cluster disappear, including apparently common and widespread haplotypes such as H and G. An alternative is that a geographically widespread genetic bottleneck occurred in *spelaea* populations sometime after 48 000 BP (the last occurrence of a haplotype external to the star radiation) which removed the earlier lineages. A subsequent re-invasion and/or range expansion of haplotype B is consistent with it being the most widespread and numerous mitochondrial sequence across the *spelaea* range (12 out of 17 specimens) after 48 000 BP. Members of the star radiation (haplotypes B and D) are first detected at 46 000 and 50 000 BP, suggesting that this group had started diversifying before the putative genetic bottleneck. The other derived haplotypes (A, C and E) may have also existed at this point, or evolved subsequently as populations containing haplotype B expanded in range and numbers. Such a pattern could be produced by the localized extinction of populations (e.g. across Eurasia), followed by the replacement with individuals from other parts of the range (e.g. eastern Beringia), as has previously been detected in Pleistocene bison and mammoth populations (Shapiro *et al.* 2004; Debruyne *et al.* 2008).

It is difficult to identify a cause for the putative bottleneck in *spelaea* populations without a precise date estimate for the event. However, it is notable that a major genetic alteration is also detectable in west Beringian bison populations between 48 000 and 50 000 BP (Cooper and Shapiro, unpublished data). This is especially significant because bison are thought to be a primary prey item for Beringian *spelaea* (Guthrie 1990; Matheus *et al.* 2003), suggesting that a population bottleneck might reflect major changes in prey availability or ecology. Alternatively, it is possible that a larger environmental change is occurring across the entire Beringian community around 48 000–50 000 BP, as this period has been identified as Heinrich event 5a (Hemming 2004), a pronounced cold stadial between Dansgaard-Oeschger (DO) interstadials 14 and 15 (GRIP 1993). Heinrich events occur during the coldest DO stadials, and are associated with catastrophic iceberg outbursts and rafting in the North Atlantic (Heinrich 1988). Heinrich events are associated with major environmental changes across the Holarctic, including neighbouring terrestrial and marine areas such as the Northwest Pacific (Gorbarenko *et al.* 2007).

With the current data, it is impossible to identify the precise nature of the genetic changes taking place in *spelaea* populations during the late Pleistocene. However, our data



add to the growing body of evidence that large-scale shifts in climate during MIS 3 were associated with major alterations in genetic structure in megafaunal populations (Barnes *et al.* 2002, 2007; Guthrie 2003; Shapiro *et al.* 2004; Hofreiter *et al.* 2007; Debruyne *et al.* 2008) well before the end-Pleistocene megafaunal extinctions in the northern hemisphere. As more data become available, integrated with detailed records of vegetative change, it should become possible to reconstruct the environment in Beringia throughout the LP to investigate the relationship between climate change and the response of megafaunal populations, providing key background information to understand their subsequent extinctions.

### Acknowledgements

Financial support was provided by Linacre College, NERC, ORADS, BBSRC, the Royal Society, and the Leverhulme Trust. Analyses in Mainz were funded in part by GEOCYCLES. We are grateful to R. Wayne, J. Leonard, and P. Matheus, H. Hemmer, G.T. Jefferson, and the George C. Page Museum for assistance and comments. We especially thank the many individuals and museums that provided access to samples, listed in the Supporting Information.

### References

- Bandelt HJ, Forster P, Rohlf A (1999) Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution*, **16**, 37–48.
- Barnes I, Matheus P, Shapiro B, Jensen D, Cooper A (2002) Dynamics of Pleistocene population extinctions in Beringian brown bears. *Science*, **295**, 2267–2270.
- Barnes I, Shapiro B, Lister AM *et al.* (2007) Genetic structure and extinction of the woolly mammoth, *Mammuthus primigenius*. *Current Biology*, **17**, 1–4.
- Barnett R, Yamaguchi N, Barnes I, Cooper A (2006) The origin, current diversity, and future conservation of the modern lion (*Panthera leo*). *Proceedings of the Royal Society B: Biological Sciences*, **273**, 2159–2168.
- Barnosky AD, Koch PL, Feranec RS, Wing SL, Shabel AB (2004) Assessing the causes of Late Pleistocene extinctions on the continents. *Science*, **306**, 70–75.
- Baryshnikov G, Boeskorov G (2001) The Pleistocene cave lion *Panthera spelaea* (Carnivora, Felidae) from Yakutia, Russia. *Cranium*, **18**, 7–24.
- Burger J, Rosendahl W, Loreille O *et al.* (2004) Molecular phylogeny of the extinct cave lion *Panthera leo spelaea*. *Molecular Phylogenetics and Evolution*, **30**, 841–849.
- Burns JA (1996) Vertebrate Palaeontology of the alleged ice-free corridor: the meat of the matter. *Quaternary International*, **32**, 107–112.
- Christiansen P (2008) Phylogeny of the great cats (Felidae: Pantherinae), and the influence of fossil taxa and missing characters. *Cladistics*, **24**, 1–16.
- Cracraft J, Feinstein J, Vaughn J, Helm-Bychowski K (1998) Sorting out tiger (*Panthera tigris*): mitochondrial sequences, nuclear inserts, systematics, and conservation genetics. *Animal Conservation*, **1**, 139–150.
- Debruyne R, Chu G, King CE *et al.* (2008) Out of America: ancient DNA evidence for a new world origin of late quaternary woolly mammoths. *Current Biology*, **18**, 1320–1326.
- Drummond AJ, Rambaut A (2007) BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology*, **7**, 214.
- García García N (2001) *Los Carnívoros de los Yacimientos Pleistocenos de la Sierra de Atapuerca*. PhD Thesis, Universidad Complutense de Madrid, Madrid, Spain.
- Gorbarenko SA, Goldberg EL, Kashgarian M *et al.* (2007) Millennium scale environmental changes of the Okhotsk Sea during last 80 kyr and their phase relationship with global climate changes. *Journal of Oceanography*, **63**, 609–623.
- GRIP (1993) Climate instability during the last interglacial period recorded in the GRIP ice core. *Nature*, **364**, 203–207.
- Groiss JT (1996) Der Höhlentiger *Panthera tigris spelaea* (Goldfuss). *Neues Jahrbuch für Geologie und Paläontologie*, **Mh. 1996/7**, 399–414.
- Guthrie RD (1990) *Frozen Fauna of the Mammoth Steppe: the Story of Blue Babe*. The University of Chicago Press, London.
- Guthrie RD (2003) Rapid body size decline in Alaskan Pleistocene horses before extinction. *Nature*, **426**, 169–171.
- Harington CR (1969) Pleistocene remains of the lion-like cat (*Panthera atrox*) from the Yukon Territory and northern Alaska. *Canadian Journal of Earth Sciences*, **6**, 1277–1288.
- Harington CR (1971) A Pleistocene lion-like cat (*Panthera atrox*) from Alberta. *Canadian Journal of Earth Sciences*, **8**, 170–174.
- Harington CR (1980) Pleistocene mammals from Lost Chicken Creek, Alaska. *Canadian Journal of Earth Sciences*, **17**, 168–198.
- Heinrich H (1988) Origin and consequences of cyclic ice rafting in the Northeast Atlantic Ocean during the past 130 000 years. *Quaternary Research*, **29**, 142–152.
- Hemming SR (2004) Heinrich events: massive late Pleistocene detritus layers of the North Atlantic and their global climate impact. *Review of Geophysics*, **42**, RG1005.
- Herrington SJ (1986) *Phylogenetic relationships of the wild cats of the world* PhD, University of Kansas.
- Ho SYW, Phillip MJ, Cooper A, Drummond AJ (2005) Time dependency of molecular rate estimates and systematic overestimation of recent divergence times. *Molecular Biology and Evolution*, **22**, 1561–1568.
- Hofreiter M, Munzel S, Conard NJ *et al.* (2007) Sudden replacement of cave bear mitochondrial DNA in the late Pleistocene. *Current Biology*, **17**, R122–R123.
- Jae-Heup K, Eizirik E, O'Brien SJ, Johnson WE (2001) Structure and patterns of sequence variation in the mitochondrial DNA control region of the great cats. *Mitochondrion*, **1**, 279–292.
- Kim J-H, Antunes A, Luo S-J *et al.* (2006) Evolutionary analysis of a large mtDNA translocation (numt) into the nuclear genome of the *Panthera* genus species. *Gene*, **366**, 292–302.
- Kurtén B (1968) *Pleistocene Mammals of Europe*. Weidenfeld and Nicolson, London.
- Kurtén B (1985) The Pleistocene lion of Beringia. *Annales Zoologici Fennici*, **22**, 177–121.
- Kurtén B, Anderson E (1980) *Pleistocene Mammals of North America*. Columbia University Press, New York.
- Lopez JV, Yuhki N, Masuda R, Modi W, O'Brien SJ (1994) Numt, a recent transfer and tandem amplification of mitochondrial-DNA to the nuclear genome of the domestic cat. *Journal of Molecular Evolution*, **39**, 174–190.
- Lopez JV, Cevario S, O'Brien SJ (1996) Complete nucleotide sequences of the domestic cat (*Felis catus*) mitochondrial genome and a transposed mtDNA tandem repeat (Numt) in the nuclear genome. *Genomics*, **33**, 229–246.

- Martin PS, Steadman DW (1999) 2. Prehistoric extinctions on islands and continents. In: *Extinctions in Near Time: Causes, Contexts, and Consequences* (ed. MacPhee RDE), pp. 17–50. Kluwer Academic/Plenum Publishing, New York.
- Matheus P, Guthrie RD, Kunz ML (2003) Predator-Prey links in Pleistocene East Beringia: evidence from stable isotopes. In: *3rd International Mammoth Conference*, Dawson City, Yukon Territory.
- Rambaut A, Drummond AJ (2007) *Tracer v1.4*. Institute of Evolutionary Biology, University of Edinburgh, Edinburgh, Scotland.
- Shapiro B, Drummond AJ, Rambaut A *et al.* (2004) Rise and fall of the Beringian steppe bison. *Science*, **306**, 1561–1565.
- Simpson GG (1941) Large Pleistocene Felines of North America. *American Museum Novitates*, **1136**, 1–27.
- Sotnikova M, Nikolskiy P (2006) Systematic position of the cave lion *Panthera spelaea* (Goldfuss) based on cranial and dental characters. *Quaternary International*, 142–143, 218–228.
- Stuart AJ (1991) Mammalian extinctions in the Late Pleistocene of Northern Eurasia and North-America. *Biological Reviews of the Cambridge Philosophical Society*, **66**, 453–562.
- Stuart AJ, Lister AM (2007) Patterns of Late Quaternary megafaunal extinctions in Europe and northern Asia. *Cour Forschung-Institute Senckenberg*, **259**, 287–297.
- Turner A, Antón M (1997) *The Big Cats and Their Fossil Relatives*. Columbia University Press, New York.
- Vereshchagin NK (1971) Cave lions of Holarctics [in Russian]. *Trudy Zoologicheskogo Instituta*, **49**, 123–199.
- Weinstock J, Willerslev E, Sher A *et al.* (2005) Evolution, systematics, and phylogeography of pleistocene horses in the new world: a molecular perspective. *PLoS Biology*, **3**, e241.
- Wilson MC (1996) Late quaternary vertebrates and the opening of the ice-free corridor, with special reference to the genus bison. *Quaternary International*, **32**, 97–105.
- Yamaguchi N, Cooper A, Werdelin L, MacDonald DW (2004) Evolution of the mane and group-living in the lion (*Panthera leo*): a review. *Journal of Zoology*, **263**, 329–342.
- 
- Ross Barnett: ancient DNA, phylogeography of felids and Pleistocene extinctions. Beth Shapiro: ancient DNA, virus evolution and phylogenetic methods. Ian Barnes: genetic changes of megafauna during the late Pleistocene. Simon Y. W. Ho: Bayesian phylogenetic methods and molecular dating. Joachim Burger: ancient DNA. Nobuyuki Yamaguchi: taxonomy and ecology of felids. Thomas F. G. Higham: radiocarbon dating and AMS methods. H. Todd Wheeler: experimental palaeontology and American Pleistocene felids. Wilfried Rosendahl: European Pleistocene cave fauna and quaternary studies. Marina Sotnikova: Pleistocene felids. Tatiana Kuznetsova: Mammoth fauna of Eurasia. Gennady F. Baryshnikov: taxonomy and evolution of carnivora and northern Eurasian Pleistocene mammals. Larry D. Martin: sabretoothed carnivores, Pleistocene felids and palaeopathology. C. Richard Harington: quaternary zoology and palaeontology. James A. Burns: Pleistocene mammalian palaeobiology and zoogeography of Alberta and North America. Alan Cooper: ancient DNA, island extinction and Pleistocene megafauna.
- 

### Supporting information

Additional supporting information may be found in the online version of this article:

**Table S1** List of all samples extracted during this study along with museum accession numbers and sample provenance. Calibrated radiocarbon dates were calculated using CalPal (Danzeglocke *et al.* 2008).

**Table S2** Primer sequences, primer combinations, annealing temperatures ( $T_a$ ), and number of thermal cycles used.

**Table S3** Number of amplifications, cloning, and replication for lion samples.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.